

JOURNAL OF DIFFERENTIAL EQUATIONS 30, 64–88 (1978)

# The Local Cauchy Problem in $\mathbb{R}^2$ at a Point Where Two Characteristic Curves Have a Common Tangent

BENT BIRKELAND

*Matematisk Institutt, Universitetet i Oslo, Postboks 1053,  
Blindern, Oslo 3, Norway*

AND

JAN PERSSON

*Institutt for matematiske realfag, Universitetet i Tromsø,  
Postboks 953, N-9001 Tromsø, Norway*

Received August 10, 1977

## 1. INTRODUCTION

We consider a linear partial differential operator in  $\mathbb{R}^2$  with principal part

$$P(x, D) = x_2^2 D_1^2 - D_2^2,$$

where  $D_1 = \partial/\partial x_1$  and  $D_2 = \partial/\partial x_2$ . The interesting feature of this operator is that its characteristics are simple when  $x_2 \neq 0$  but double when  $x_2 = 0$ . The characteristic curves through the point  $(c, 0)$  are the parabolas

$$x_1 = \pm x_2^2/2 + c.$$

They have a common tangent at  $(c, 0)$ . We are interested in the Cauchy problem for such operators when data are given on a curve characteristic at the origin.

The very first published results on this problem seem to be the following two theorems by Treves [18] concerning the operator

$$P_\lambda(x, D) = x_2^2 D_1^2 - D_2^2 - \lambda D_1, \quad (1.1)$$

where  $\lambda$  is a real parameter.

**THEOREM 1.1** (Treves [18, Theorem II]). *If  $\lambda$  is an odd positive integer then there exist  $C^\infty$ -functions  $u$  in  $\mathbb{R}^2$  such that*

$$P_\lambda u = 0, \quad \text{supp } u = \{x; x_1 \geq x_2^2/2\}.$$

**THEOREM 1.2** (Treves [18, Theorem I]). *Suppose that  $\lambda$  in (1.1) is real but not an odd positive integer. Let  $\Omega \subset \mathbb{R}^2$  be any open set, and  $F \subset \Omega$  a closed subset such that for some real  $a$  the set*

$$K = \{x; x \in \Omega, x_1 \leq a\} \cap F$$

*is compact. Then there exist an integer  $m \geq 2$  depending solely on  $\lambda$  and a neighborhood  $U$  of  $K$  such that any function  $u \in C^m(\Omega)$  satisfying*

$$P_\lambda u = 0, \quad \text{supp } u \subset F \quad (1.2)$$

*must vanish in  $U$ .*

Treves remarks that it seems likely that  $m$  can be chosen equal to 2 for all  $\lambda$  [18, footnote, p. 230]. We prove a stronger result.

**THEOREM 1.3.** *Let  $P_\lambda$ ,  $\Omega$ ,  $F$ , and  $K$  be as in the hypothesis of Theorem 1.2. Then there is an open set  $U \supset K$  such that any  $u \in \mathcal{D}'(\Omega)$  fulfilling (1.2) must vanish in  $U$ .*

The proof is given in Section 2. We use Schwartz' structure theorem for distributions. This idea has been used earlier in proofs of uniqueness theorems (see Persson [8]). In [9] Persson conjectures that uniqueness cones can always be used to decide whether there is uniqueness in the local Cauchy problem when the coefficients are analytic. Theorem 1.3 shows that this conjecture is false.

Theorem 1.1 gives us null solutions with support in  $\{x; x_1 \geq x_2^2/2\}$  when  $\lambda$  is an odd positive integer. Theorem 1.4 below shows that in a somewhat modified solution space we have "null solutions" for all real  $\lambda$ .

**THEOREM 1.4.** *There exist continuous functions  $u \neq 0$  defined on  $\mathbb{R}$  with values in the space  $H'(\mathbb{C})$  of analytic functionals over  $\mathbb{C}$  such that with  $P_\lambda u$  defined in the natural way*

$$u(x_1) = 0, \quad x_1 \leq 0, \quad P_\lambda u = 0, \quad (x_1, x_2) \in \mathbb{R} \times \mathbb{C}.$$

*Moreover for each  $x_1 \geq 0$  the functional  $u(x_1)$  is carried by the set*

$$\{x_2; x_2 \in \mathbb{C}, |x_2| \leq (2x_1)^{1/2}\}.$$

As to the definition of analytic functionals and elementary facts about them we refer the reader to Treves [17, Chap. 9]. From Theorem 1.3 we see that  $u$  in Theorem 1.4 cannot be a distribution unless  $\lambda$  is an odd positive integer. We prove Theorem 1.4 in Section 3.

The next problem which we consider is whether there exist nontrivial solutions of  $P_\lambda(D)u = 0$  with supports to the right of the leftmost characteristic curve

through the origin. It turns out that such solutions always exist at least locally. We prove a theorem for a more general situation giving this result as a special case. At first we supplement our notation by letting  $|x| = \max(|x_1|, |x_2|)$ ,  $x \in \mathbb{R}^2$ . For  $\xi = (\xi_1, \xi_2) \in \mathbb{Z}^2$ ,  $\xi_j \geq 0$ , and  $D = (D_1, D_2) = (\partial/\partial x_1, \partial/\partial x_2)$ , we let  $D^\xi = D_1^{\xi_1} D_2^{\xi_2}$ . For  $d = (d_1, d_2) \in \mathbb{R}^2$ ,  $d_1 \geq 1$ ,  $d_2 \geq 1$ , we let  $\xi d = \xi_1 d_1 + \xi_2 d_2$ . We also let  $|\xi| = \xi_1 + \xi_2$  and  $\xi d^{\xi d - 1} = 1$ ,  $\xi d = 0$ , and  $(\xi d - 2)^{\xi d - 3} = 1$ ,  $0 \leq \xi d - 2 \leq 1$ .

We notice that our restrictions on  $\xi$  and  $d$  imply that  $\xi \rightarrow \xi d^{\xi d - 1}$  and  $d \rightarrow \xi d^{\xi d - 1}$  both are nondecreasing.

Now we are ready to state our last theorem.

**THEOREM 1.5.** *Let  $\epsilon > 0$  and let  $\Omega = \{x; x \in \mathbb{R}^2, |x| < \epsilon\}$ . Let  $a_\alpha \in C^\infty(\Omega)$ ,  $|\alpha| \leq 1$  and  $b(x_2) \in C^\infty(\{x_2; |x_2| < \epsilon\})$ . Let  $m_\alpha \geq 0$ ,  $|\alpha| \leq 1$ ,  $m > 0$ , and  $r, 0 < r \leq 1$ , and  $d_1 = d_2$ ,  $1 < d_1 \leq 2$  be constants. We assume that with  $d = (d_1, d_2)$*

$$|D^\xi a_\alpha(x)| \leq m_\alpha r^{-|\xi|} \xi d^{\xi d - 1}, \quad |\alpha| \leq 1, \quad \text{all } \xi, x \in \Omega, \quad (1.3)$$

and

$$|D_2^j b(x_2)| \leq m r^{-|j|} \xi d^{\xi d - 1}, \quad \text{all } j \geq 0, \quad |x_2| < \epsilon. \quad (1.4)$$

Let

$$\Psi(t) = \int_0^t \min(0, b(s)) ds, \quad 0 \leq t < \epsilon,$$

and

$$\Psi(t) = \int_0^t \max(0, b(s)) ds, \quad 0 < -t < \epsilon.$$

Let

$$P(x, D) = D_2(b(x_2) D_1 + D_2) - \sum_{|\alpha| \leq 1} a_\alpha(x) D^\alpha, \quad x \in \Omega. \quad (1.5)$$

Then there exist a neighborhood  $\Omega_0$  of 0 and a  $u \in C^\infty(\Omega_0)$  such that

$$P(x, D)u = 0 \text{ in } \Omega_0 \quad \text{and} \quad 0 \in \text{supp } u \subset \{x; x_1 \geq \Psi(x_2)\}.$$

*Remark 1.* If  $b(0) \neq 0$ , then near the origin we may use new coordinates with the characteristic curves as coordinate axes. We may choose this system such that in this system

$$P(x, D) = D_1 D_2 + a_1 D_1 + a_2 D_2 + a_0.$$

Then we solve the Goursat problem,

$$P(x, D)u = 0, \quad u(x_1, 0) = \varphi(x_1), \quad u(0, x_2) = 0.$$

Here we choose  $\varphi(x_1) = 0$ ,  $x_1 \leq 0$ ,  $\varphi(x_1) > 0$ ,  $x_1 > 0$ ,  $\varphi \in C^\infty(\mathbb{R})$ . The corresponding solution  $u$  (see, for instance, Persson [7, proof of Theorem 2]) has  $\text{supp } u \subset \{x; x_1 \geq 0\}$  and  $0 \in \text{supp } u$ . This is the idea behind Goursat's<sup>1</sup> original construction of null solutions with data given on a characteristic line of the wave operator. It is not hard to show that  $u \in C^\infty$  too. Then we transform back and there we have our wanted solution.

*Remark 2.* The characteristic curves of (1.5) through  $(c, 0)$  are the line  $x_1 = c$  and  $x_1 = c + \varphi(x_2)$ , where

$$\varphi(x_2) = \int_0^{x_2} b(t) dt. \quad (1.6)$$

They have a common tangent at  $(c, 0)$  precisely when  $\varphi'(0) = b(0) = 0$ .

*Remark 3.* After obvious modifications the theorem is also valid for operators which can be transformed into (1.5) by suitable coordinate changes. We mention two such operators:

(a) The operator  $P_\lambda$  of (1.1) is transformed into (1.5) by the coordinate shift

$$x'_1 = x_1 + x_2^2/2, \quad x'_2 = x_2.$$

(b) If we allow the function  $b$  in (1.5) to be of the form  $b(x) = b_1(x_1) b_2(x_2)$  with both  $b_j$  satisfying estimates of the form (1.4) the resulting operator is transformed back to (1.5) by

$$x'_1 = \int_0^{x_1} (b_1(s))^{-1} ds, \quad x'_2 = x_2.$$

Then it can be shown that the coefficients of the lower order still satisfy estimates of (1.4) type.

The proof of Theorem 1.5 is given in Section 4. Some soft and some harder auxiliary results are proved in Sections 5 and 6.

How then is Theorem 1.5 related to known results on nonuniqueness? The first construction of a null solution seems to be the one already cited by Goursat. In the constant coefficient case we refer to Tihonov [19], Täcklind [16], Hörmander [3, Theorem 3.2] or [4, Theorem 5.2.2, p. 121], and Persson [12]. In the case of analytic coefficients we refer to [4, Theorem 5.2.1] when the initial hypersurface is simply characteristic and to Persson [10, 11, 13] when the multiplicity of the initial hypersurface is arbitrary but constant. Later Bronšteĭn [2] extended the results in [10] to nonlinear problems. Komatsu [6] has also constructed null solutions by another method. In all the literature cited above

<sup>1</sup> E. Goursat, "Leçons sur l'intégration des équations aux dérivées partielles de second ordre," Tome II, pp. 303–308, Hermann, Paris, 1898.

the initial characteristic hypersurface has constant multiplicity. If we let the data of  $P_\lambda(D)u = 0$  be given on  $x_1 = -x_2^2/2$  with  $P_\lambda$  from (1.1) then the multiplicity of the initial curves is 2 at  $x = 0$  and 1 for  $x \neq 0$ . So this case is not contained in the results cited above. We allow the coefficients to be in non-analytic Gevrey classes in Theorem 1.5. In [10] it is indicated how one may weaken the hypothesis in this direction when the multiplicity of the characteristic initial surface is constant.

If the principal part vanishes on the initial hypersurface, Hörmander [5, Theorem 2.2] has given some examples of null solutions when the coefficients are analytic. We do not intend to give any complete survey of results on uniqueness and nonuniqueness in the characteristic Cauchy problem but would like to cite Baouendi and Goulaouic [1]. They have characterized other types of characteristic Cauchy problems where one cannot construct  $C^\infty$ -null solutions.

*Post Scriptum.* One can also see the proof of Theorem 1.5 as a construction of a solution of the Cauchy problem with zero initial data given on  $x_2 = 0$ . Here one enters into a long range of results. At the time of the writing we found no results giving room for one characteristic curve to oscillate around the other one as in Theorem 1.5. However some results are more general in other aspects. Then after the paper was completed we learned from Zentralblatt about Ivrii [E7] and his striking results on the Cauchy problem for operators with hyperbolic principal part. He treats the case when coefficients and data are in Gevrey classes. His results cover our result in the proof of Theorem 1.5. We still think that our more direct construction and our point of view justify the publication of our paper. We have included some papers on the Cauchy problem in the  $x_2$ -direction including that by Ivrii in an extra reference list following the ordinary one.

It also happened that the first author tried to compute the best constant  $c$  in Lemma 5.1. He conjectured that  $c = 4$  is the best one. Then Arne Strøm, Oslo, and Robert Fossum and Erik Sparre Andersen, Copenhagen, showed us how to prove this fact which goes back to Abel. We thank all these people. Section 5 is rewritten accordingly.

## 2. PROOF OF THEOREM 1.3

Let  $K \subset \mathbb{R}^2$  and  $\epsilon > 0$ ,  $c \in \mathbb{R}$ . We define

$$K_\epsilon = \{x; \text{dist}(x, K) \leq \epsilon\}$$

and

$$\Omega_c = \{x; x_1 < c\}.$$

Now let the sets  $\Omega$ ,  $K$ ,  $F$ , and the number  $a$  be as in Theorems 1.2 and 1.3. We

look at a distribution solution  $u$  of  $P_\lambda u = 0$  in  $\Omega$  with  $\text{supp } u \subset F$ . We want to show that  $u = 0$  in some neighborhood of  $K$ .

We start by choosing  $\eta > 0$  so small that

$$K_{3\eta} \subset \Omega.$$

It follows that the closure of the set

$$(K_{3\eta} - K_\eta) \cap \{x; x_1 = a\}$$

is compact and disjoint from  $F$ . Thus it has a positive distance to  $F$ . Therefore we can choose a real number  $c$  such that  $a < c < a + \eta$ , and such that

$$K_\eta \cap F \cap \Omega_c = K_{3\eta} \cap F \cap \Omega_c. \quad (2.1)$$

Now we use cut-off functions to split  $u$  into a sum

$$u = u_1 + u_2,$$

where

$$\text{supp } u_1 \subset K_{2\eta} \cap F, \quad (2.2)$$

and

$$\text{supp } u_2 \cap K_\eta = \emptyset, \quad \text{supp } u_2 \subset F.$$

From (2.1) it then follows that

$$\text{supp } u_2 \cap K_{3\eta} \cap \Omega_c = \emptyset. \quad (2.4)$$

Since

$$\text{supp } P_\lambda u_1 \subset (\text{supp } u_1) \cap (\text{supp } u_2),$$

we also have

$$\text{supp } P_\lambda u_1 \cap \Omega_c = \emptyset.$$

We extend  $u_1$  by letting it be zero outside  $K_{3\eta}$  to obtain  $u_1 \in \mathcal{D}'(\mathbb{R}^2)$  and

$$P_\lambda u_1 = 0, \quad x \in \Omega_c. \quad (2.5)$$

It follows from Schwartz's theorem on the structure of distributions with compact support [14, Théorème 26, p. 91] and (2.2) that there exist a positive integer  $m$  and continuous functions  $f_\alpha$ ,  $|\alpha| \leq m$  with  $\text{supp } f_\alpha \subset K_{3\eta}$  such that

$$u_1 = \sum_{|\alpha| \leq m} D^\alpha f_\alpha.$$

To simplify notations we choose a real number  $b$  such that

$$K_{4\eta} \subset \{x; x_1 > b, x_2 > b\}. \quad (2.6)$$

For continuous functions  $g$  with  $\text{supp } g \subset \{x; x_1 > b, x_2 > b\}$  we define

$$D_1^{-1}g(x) = \int_b^{x_1} g(s, x_2) ds, \quad D_2^{-1}g(x) = \int_b^{x_2} g(x_1, t) dt.$$

It is clear that  $D_1^{-1}g$  and  $D_2^{-1}g$  vanish when  $x_1 \leq b$  or  $x_2 \leq b$ . It is obvious that  $D_1, D_2, D_1^{-1}, D_2^{-1}$  all commute and that

$$D_j D_j^{-1}g = D_j^{-1} D_j g = g, \quad j = 1, 2.$$

Then we let  $D_j^{-n} = (D_j^{-1})^n, n \geq 0$ . We now see that we may write

$$u_1 = D_1^m D_2^m f$$

with

$$f = \sum_{|\alpha| \leq m} D_1^{\alpha_1 - m} D_2^{\alpha_2 - m} f_\alpha.$$

We also notice that (2.4) and  $\text{supp } f_\alpha \subset K_{3\eta} \mid \alpha \mid \leq m$ , imply that

$$\text{supp } f \subset \{x; x_1 > b + \eta, x_2 > b + \eta\}. \quad (2.7)$$

Now we regularize in the  $x_1$ -direction. Let  $\varphi \in C_0^\infty(\mathbb{R}^1)$  satisfy  $\int \varphi = 1$ ,  $\varphi(t) = 0$  for  $|t| > \frac{1}{2}$ . Then we let

$$\varphi_\epsilon(x_1) = \epsilon^{-1} \varphi(x_1/\epsilon)$$

and

$$v = v_\epsilon = u_1^{*'} \varphi_\epsilon,$$

where  $^{*'}$  denotes convolution in the  $x_1$ -variable. The coefficients of  $P_\lambda$  do not depend on  $x_1$  so we have  $P_\lambda v = (P_\lambda u_1)^{*'} \varphi_\epsilon$ . Hence (2.5) gives us

$$P_\lambda v = 0 \quad \text{in } \Omega_{c-\epsilon}. \quad (2.8)$$

We choose  $\epsilon$  such that

$$0 < \epsilon < c - a < \eta.$$

We notice that this and (2.2) gives us

$$\text{supp } v \subset K_{4\eta} \subset \{x; x_1 > b, x_2 > b\}.$$

So we have

$$v(x) = (D_1^m D_2^m f)^{*'} \varphi_\epsilon = D_2^m (f^{*'} D_1^m \varphi_\epsilon) = D_2^m g,$$

where

$$g = f^{*'} D_1^m \varphi_\epsilon$$

is a continuous function smooth in  $x_1$ .

Now we rewrite (2.8) as

$$D_2^2 D_2^m g = x_2^2 D_1^2 D_2^m g - \lambda D_1 D_2^m g, \quad x \in \Omega_{c-\epsilon}. \quad (2.9)$$

We notice that  $\text{supp } g \subset \{x; x_1 > b, x_2 > b\}$ . We also notice that for  $m \geq 2$

$$\begin{aligned} D_2^m (x_2^2 D_1^2 g) &= \sum_{j=0}^2 \binom{m}{j} D_2^j (x_2)^2 D_2^{m-j} D_1^2 g \\ &= x^2 D_2^m D_1^2 g + 2m x_2 D_2^{m-1} D_1^2 g + m(m-1) D_2^{m-2} D_1^2 g. \end{aligned}$$

We also have

$$\begin{aligned} D_2^{m-1} (2m x_2 D_1^2 g) &= m \sum_{j=1}^1 \binom{m-1}{j} D_2^j (2x_2) D_2^{m-1-j} D_1^2 g \\ &= 2m x_2 D_2^{m-1} D_1^2 g + m(m-1) 2 D_2^{m-2} D_1^2 g. \end{aligned}$$

We put these things together and get

$$\begin{aligned} D_2^{2+m} g &= D_2^m (x_2^2 D_1^2 g) - D_2^{m-1} (2m x_2 D_1^2 g) \\ &\quad + D_2^{m-2} (m(m-1) D_1^2 g) - D_2^m (\lambda D_1 g) \\ &= D_2^m [x_2 D_1^2 g - D_2^{-1} (2m x_2 D_1^2 g) + D_2^{-2} (m(m-1) D_1^2 g) \\ &\quad - \lambda D_1^2 g] = D_2^m h = D_2^{m+2} D_2^{-2} h. \end{aligned}$$

We see that  $h$  is continuous. It is smooth in the  $x_1$ -variable and

$$\text{supp } h \subset \{x; x_1 > b, x_2 > b\}.$$

This shows that

$$g = D_2^{-2} h, \quad x \in \Omega_{c-\epsilon}. \quad (2.10)$$

For  $m = 1$  it is still simpler. For  $m = 0$  it is obvious. Now assume that  $m$  is the smallest integer, positive or not, such that for some continuous  $g$  with  $\text{supp } g \subset \{x; x_1 > b, x_2 > b\}$

$$v = D_2^m g \quad \text{in } \Omega_{c-\epsilon},$$

$g$  being smooth in  $x_1$ .

The calculation above for  $m \geq 0$  and an obvious argument for  $m < 0$  show that (2.10) is true for some  $h$  fulfilling the same regularity condition as  $g$ . So  $m$  was not minimal and  $v$  restricted to  $\Omega_{c-\epsilon}$  is in  $C^\infty(\Omega_{c-\epsilon})$ . Now  $v_\epsilon$  satisfies all the conditions of Theorem 1.2 in  $\Omega_{c-\epsilon}$  so  $v_\epsilon = 0$  in  $\Omega_{c-\epsilon}$ . That means that  $u_1 = 0$  in  $\Omega_c$  since  $v_\epsilon \rightarrow u_1$  there when  $\epsilon \rightarrow 0$ . Then (2.4) implies that  $u_1 = u$  in  $\tilde{K}_{3\eta} \cap \Omega_c$ . Let  $U = \tilde{K}_{2\eta} \cap \Omega_c$ . The theorem is proved.



## 3. PROOF OF THEOREM 1.4

In this proof we prefer to abandon the multi-index notation and denote points in  $\mathbb{R}^2$  by  $(x, y)$  instead of  $(x_1, x_2)$ . We also use  $D_x = \partial/\partial x$  and  $D_y = \partial/\partial y$ . Our equation  $P_\lambda u = 0$  then reads

$$(y^2 D_x^2 - D_y^2 - \lambda D_x) u(x, y) = 0. \quad (3.1)$$

The Fourier-Borel transform with respect to  $y$  transforms this into

$$(D_z^2 D_x^2 - z^2 - \lambda D_x) w(x, z) = 0. \quad (3.2)$$

We are interested in solutions  $w(x, z)$  which are continuous functions of  $(x, z) \in \mathbb{R} \times \mathbb{C}$ , analytic in  $z$  for fixed  $x$  and vanishing for  $x < 0$ . We seek solutions of the form

$$w(x, z) = \sum_{j=0}^{\infty} \varphi_j(x) z^j / j!, \quad (3.3)$$

where the  $\varphi_j$  are in  $C^2(\mathbb{R})$  vanishing for  $x \leq 0$ . Formal substitution of (3.3) into (3.2) gives us the following differential equations for  $\varphi_j$ :

$$\varphi_2'' = \lambda \varphi_0', \quad \varphi_3'' = \lambda \varphi_1',$$

and

$$\varphi_{j+2}'' = \lambda \varphi_j' + j(j-1) \varphi_{j-2}, \quad j \geq 2.$$

We notice that  $\varphi_j \in C^2(\mathbb{R})$ ,  $\varphi_j(x) = 0$ ,  $x \leq 0$  implies that  $\varphi_j(0) = \varphi_j'(0) = 0$ . We define

$$D^{-1}g(x) = \int_0^x g(t) dt.$$

All this implies that

$$\varphi_2 = \lambda D^{-1} \varphi_0, \quad \varphi_3 = \lambda D^{-1} \varphi_1,$$

and

$$\varphi_{j+2} = \lambda D^{-1} \varphi_j + j(j-1) D^{-2} \varphi_{j-2}, \quad j \geq 2.$$

Thus all  $\varphi_j$  can be expressed in terms of the two first ones as follows

$$\begin{aligned} \varphi_{2j} &= \theta(0, \lambda, j) D^{-j} \varphi_0, \\ \varphi_{2j+1} &= \theta(1, \lambda, j) D^{-j} \varphi_1, \quad j \geq 1, \end{aligned}$$

where  $\theta(i, \lambda, j)$  are complex numbers fulfilling the recursive formulas

$$\theta(0, \lambda, j+1) = \lambda \theta(0, \lambda, j) + 2j(2j-1) \theta(0, \lambda, j-1), \quad j \geq 0,$$

and

$$\theta(1, \lambda, j+1) = \lambda \theta(1, \lambda, j) + (2j+1) 2j \theta(1, \lambda, j-1), \quad j \geq 0$$

with

$$\theta(i, \lambda, 0) = 1, \theta(i, \lambda, j) = 0, \quad j < 0.$$

It follows by induction that

$$|\theta(0, \lambda, j)| \leq \prod_{k=1}^j (2k + |\lambda|).$$

Since

$$D^{-j} \varphi_0(x) = ((j-1)!)^{-1} \int_0^x \varphi_0(t) (x-t)^{j-1} dt, \quad j \geq 1,$$

a simple computation shows that the series

$$w_0(x, z) = \sum_{j=1}^{\infty} \varphi_{2j} z^{2j} / (2j)!$$

converges in  $\mathbb{R} \times \mathbb{C}$ , uniformly on compact sets, and that for fixed  $x \geq 0$  the function  $z \rightarrow w_0(x, z)$  is entire and that

$$|w_0(x, z)| \leq C_\epsilon \exp(|z|((2x)^{1/2} + \epsilon))$$

for every  $\epsilon > 0$  and

$$w_0(x, z) = 0, \quad x < 0.$$

It is also seen that  $w_0(x, z)$  solves (3.2). Quite similar statements hold for

$$w_1(x, z) = \sum_{j=0}^{\infty} \varphi_{2j+1}(x) z^{2j+1} / (2j+1)!$$

We see that for any choice of  $\varphi_0$  and  $\varphi_1$  vanishing for  $x \leq 0$  we obtain a solution  $w(x, z) = w_0(x, z) + w_1(x, z)$  of (3.2) in  $C^2(\mathbb{R} \times \mathbb{C})$ , entire of exponential growth in  $z$  for each fixed  $x$ . When we take the inverse Fourier-Borel transform of  $w(x, z)$  in the  $z$ -variable (see [17, Theorem 9.1, p. 474]), then we obtain a function  $u \in C^2(\mathbb{R}, H'(\mathbb{C}))$  solving (3.1). We also have that  $u \neq 0$  and that for fixed  $x > 0$ ,

$$\{y; y \in \mathbb{C}, |y| \leq (2x)^{1/2}\}$$

is a carrier of  $u(x)$ . The theorem is proved.

## 4. PROOF OF THEOREM 1.5

The starting point for the proof is the observation that the differential operator

$$P_2(x, D) = D_2(b(x_2)D_1 + D_2) \quad (4.1)$$

has a right inverse  $T$  which is given explicitly as the integral operator

$$Tg(x) = \int_0^{x_2} \int_0^t g(x_1 - \varphi(x_2) + \varphi(t), s) ds dt, \quad (4.2)$$

where  $\varphi' = b$  and  $\varphi(0) = 0$ . We shall use this fact when we construct null solutions of the full equation  $P(x, D)u = 0$  by successive approximations. In order to prove the convergence of these approximations we need some inequalities. They are proved in Sections 5 and 6, but we shall state them and use them in this section.

LEMMA 4.1. *Let  $\Omega \subset \mathbb{R}^2$  be open and let  $f, g \in C^\infty(\Omega)$ . Let  $m, m', r, d_1 \geq 1$ ,  $d_2 \geq 1$  be positive constants and let  $q(x) \geq 0$  in  $\Omega$  be such that with  $d = (d_1, d_2)$*

$$|D^\xi f| \leq mr^{-|\xi|} \xi d^{\xi d-1}, \quad x \in \Omega, \text{ all } \xi, \quad (4.3)$$

and

$$|D^\xi g| \leq m'r^{-|\xi|} \xi d^{\xi d-1} \exp[(1 + \xi d) q(x)], \quad x \in \Omega, \text{ all } \xi. \quad (4.4)$$

Then there exists a constant  $c$ , independent of all quantities mentioned above, such that

$$|D^\xi(fg)| \leq cmm'r^{-|\xi|} \xi d^{\xi d-1} \exp[(1 + \xi d) q(x)], \quad x \in \Omega, \text{ all } \xi. \quad (4.5)$$

COROLLARY 4.2. *Let  $f$  be as above, and let  $k$  be a positive integer. Then*

$$|D^\xi(f^{k+1})| \leq m(mc)^k r^{-|\xi|} \xi d^{\xi d-1}. \quad (4.6)$$

*Proof.* Let  $q = 0$  in the lemma and use induction in  $k$ .

LEMMA 4.3. *Let  $\Omega = \{x; x \in \mathbb{R}^2, |x_2| < \rho\}$  for some  $\rho > 0$ . Then (4.2) defines a function  $Tg \in C^\infty(\Omega)$  if  $g \in C^\infty(\Omega)$  and  $\varphi \in C^\infty(\mathbb{R})$ . Let  $g$  satisfy (4.4) with  $r, 0 < r \leq 1$ , and  $\Omega$  as above, and*

$$q(x) = e^2 |x_2|/r, \quad \rho \leq re^{-2}. \quad (4.7)$$

Let  $\varphi \in C^\infty(\{t, |t| < \rho\})$  fulfill

$$|D^{j+1}\varphi(t)| \leq mR^{-j}(jd_2)^{j\bar{d}_2-1}, \quad |t| < \rho, j = 0, 1, \dots, \quad (4.8)$$

where  $0 < r/R < \frac{1}{4}$ , and  $mc < \frac{1}{4}$ . Here  $c$  is taken from Lemma 4.1. Then with  $T$  defined in (4.2) we have

$$|D^\epsilon Tg| \leq 4m'r^{2-|\epsilon|} (\xi d - 2)^{\epsilon d - 3} \exp[(\xi d - 1)q(x)], \quad x \in \Omega, |\xi| \geq 2, \quad (4.9)$$

and

$$|D^\epsilon Tg| \leq 4m', \quad x \in \Omega, \quad |\xi| \leq 1. \quad (4.10)$$

We like to work with an operator defined in  $\Omega = \{x; x \in \mathbb{R}^2, |x_2| < \rho\}$  such that (4.8) is fulfilled and such that (1.3) and (1.4) are fulfilled with this  $\Omega$  and with  $\epsilon = \rho$ . In addition we want to have

$$\sum m_\alpha \leq (8c)^{-1}, \quad |\alpha| \leq 1. \quad (4.11)$$

Here  $c$  was introduced in Lemma 4.1. The constants  $m_\alpha$  come from (1.3). We also require that

$$|D_2^j b(x_2)| \leq mR^{-j}(jd_2)^{jd_2-1}, \quad j \geq 0, |x_2| < \rho, \quad (4.12)$$

where

$$0 < r/R < \frac{1}{4}, \quad mc < \frac{1}{4}. \quad (4.13)$$

We begin the proof by showing how the general case can be reduced to this one. We may assume that we have adjusted  $r$  and  $R$  such that the first inequality of (4.13) is satisfied and such that (1.3) is true with the new  $r$  and that (1.4) is true with  $r$  replaced by  $R$ . We define  $q$  by (4.7) and choose  $\rho$  such that  $0 < \rho < \min(re^{-2}, \epsilon)$ . Then we choose a cut-off function  $h \in C^\infty(\mathbb{R})$  such that

$$0 \leq h(x_1) \leq 1, \quad h(x_1) = 1, \quad |x_1| \leq \rho/2, \quad h(x_1) = 0, \quad |x_1| > \rho,$$

also fulfilling

$$|D^j h| \leq m'r^{-j}(jd_1)^{jd_1}, \quad x_1 \in \mathbb{R}.$$

The existence of such a function follows from [4, Lemma 5.7.1, p. 146]. Now we define

$$\begin{aligned} a'_\alpha(x) &= h(x_1) a_\alpha(x), & |x_1| \leq \rho, & |x_2| < \rho, \\ a'_\alpha(x) &= 0, & |x_1| > \rho, & |x_2| < \rho. \end{aligned}$$

Then Lemma 4.1 with  $q(x) = 0$  shows that the estimates of (1.3) are still true if we replace  $a_\alpha$  by  $a'_\alpha$ . The only change is that we may have to change the values of the constants  $m_\alpha$ . If we replace the coefficients  $a_\alpha$  in  $P(x, D)$  by  $a'_\alpha$  then we get a new operator  $P'(x, D)$  defined in  $\Omega = \{x; |x_2| < \rho, x \in \mathbb{R}^2\}$ . In a neighborhood of  $x = 0$  the equations  $P(x, D)u = 0$  and  $P'(x, D)u = 0$  have the same solutions. From now on we work with  $P'(x, D)$  and delete the primes.

If in our original coordinate system, (4.11) or the last inequality of (4.13) are not true then we choose

$$x'_1 = x_1 \quad \text{and} \quad x'_2 = tx_2,$$

with some constant  $t \geq 1$ . So we have

$$D_2 u(x) = t(D'_2 u')(x'),$$

where  $u(x) = u'(x')$ . Now  $P(x, D)u = 0$  is transformed into the equivalent equation

$$\begin{aligned} P'(x', D') u' &= D'_2(t^{-1}b(x_2) D'_1 + D'_2) u' \\ &- \sum_{|\alpha| \leq 1} t^{\alpha_2-2} a_\alpha(x) D'^\alpha u' = D'_2(b'(x'_2) D'_1 + D'_2) u' \\ &- \sum_{|\alpha| \leq 1} a'_\alpha D'^\alpha u' = 0. \end{aligned}$$

We have

$$|D'^\xi a'_\alpha(x')| \leq m_\alpha t^{\alpha_2-2-\xi_2} r^{-|\xi|} \xi d^{\xi d-1}, \quad |x'_2| < t\rho, x'_1 \in \mathbb{R}, \text{ all } \xi,$$

and

$$|D_2^j b'(x'_2)| \leq m t^{-1-j} R^{-j} (j d_2)^{j d_2-1}, \quad |x'_2| < t\rho, x' \in \mathbb{R}, j \geq 0.$$

It is now clear that with a proper choice of  $t \geq 1$  and after deleting the primes we may assume that both (4.11) and the last inequality of (4.13) are fulfilled. So we assume that this is true from the beginning.

We notice that with  $T$  from (4.2)

$$D_2(bD_1 + D_2) Tg = g, \quad g \in C^\infty(\Omega), \quad (4.14)$$

and that

$$Tg(x_1, 0) = 0, \quad x_1 \in \mathbb{R}. \quad (4.15)$$

Let  $g \in C^\infty(\Omega)$  be such that  $g(x) = 0$  in  $M = \{x; x \in \Omega, x_1 < \psi(x_2)\}$  with  $\psi$  defined in the hypothesis of Theorem 1.5. Then we assert that  $Tg(x) = 0$  in  $M$ . Let  $x \in M$ ,  $x_2 < 0$ . In  $T$  we have  $x_2 \leq t \leq 0$ ,  $t \leq s \leq 0$ . We notice that  $d/dt(\psi - \varphi) = \max(0, b) - b \geq 0$ ,  $t < 0$ , and that

$$\begin{aligned} x_1 - \varphi(x_2) + \varphi(t) - \psi(s) &< \psi(x_2) - \varphi(x_2) + \varphi(t) - \psi(t) \\ &\leq \psi(x_2) - \psi(t) + \varphi(t) - \varphi(x_2) \leq 0. \end{aligned}$$

The case  $x_2 > 0$  is now also obvious.

Now we construct a solution  $u$  of  $P(x, D)u = 0$ . We start by choosing a function  $u^0(x)$  of the form

$$u^0(x) = h(x_1), \quad x \in \Omega. \quad (4.16)$$

We choose  $h(x_1)$  such that

$$h(x_1) = 0, \quad x_1 \leq 0, \quad h(x_1) > 0, \quad x_1 > 0, \quad (4.17)$$

and such that for some  $m'' > 0$

$$|D^j h| \leq m'' R^{-j} d_1^{-1}, \quad x_1 \in \mathbb{R}, \text{ all } j \geq 0. \quad (4.18)$$

Here  $d_1$  is chosen from the hypothesis. The number  $R$  is the constant chosen below formula (4.13). We again refer to [4, Lemma 5.7.1, p. 146]. Then we define

$$f^0(x) = -P(x, D)u^0, \quad x \in \Omega, \quad (4.19)$$

and recursively for  $p \geq 1$

$$u^p(x) = T f^{p-1}(x), \quad x \in \Omega, \quad (4.20)$$

and

$$f^p(x) = \sum a_\alpha D^\alpha u^p, \quad x \in \Omega. \quad (4.21)$$

We prove that for every  $\Omega$  the series

$$\sum_p D^\xi u^p \quad (4.22)$$

converges absolutely uniformly on  $\Omega$  for all  $\xi$ . This means that  $u = \sum u^p$  is a well defined function in  $C^\infty(\Omega)$ . Now (4.16) and (4.17) tell us that  $u^0(x_1, 0) > 0$ ,  $x_1 > 0$ , while (4.15) tells us that  $u^p(x_1, 0) = 0$ ,  $x_1 > 0$ , for all  $p \geq 1$ . That shows us that

$$u(x_1, 0) > 0, \quad x_1 > 0.$$

We have

$$\text{supp } u^0 \subset \{x; x_1 \geq 0\} \subset \{x; x_1 \geq \psi(x_2)\}.$$

The argument after (4.15) then implies that

$$\text{supp } u^p \subset \{x; x_1 \geq \psi(x_2)\}.$$

So the same is true for  $u$  itself. Finally we have by (4.22), (4.20), (4.14), and (4.21) that

$$\begin{aligned} P(x, D)u &= P(x, D)u^0 + \sum_1^\infty \left( D_2(bD_1 + D_2)u^p - \sum_{|\alpha| \leq 1} a^\alpha D^\alpha u^p \right) \\ &= f^0 + \sum_1^\infty f^{p-1} - \sum_1^\infty f^p = 0. \end{aligned}$$

It remains to prove (4.22).

It follows from (4.16) and (4.18) that

$$|D^\xi u^0| \leq m'' R^{-|\xi|} \xi d^{\xi d-1}, \quad x \in \Omega, \text{ all } \xi.$$

A short calculation based on (4.18), Lemma 4.1, (1.3), and (4.12) shows that

$$|D^\xi f^0| \leq m' r^{-|\xi|} \xi d^{\xi d-1}, \quad x \in \Omega, \text{ all } \xi, \quad (4.23)$$

for some constant  $m'$ . We want to prove that

$$|D^\xi f^p| \leq 2^{-p} m' r^{-|\xi|} \xi d^{\xi d-1} \exp[(1 + \xi d) q(x)], \quad p \geq 0, x \in \Omega, \text{ all } \xi, \quad (4.24)$$

where  $q(x)$  is still given by (4.7). Now (4.23) shows that (4.24) is true for  $p = 0$ . So we assume that (4.24) has been established for some  $p \geq 0$ . Then (4.20), (4.24), and Lemma 4.3 show that

$$|D^\xi u^{p+1}| \leq 2^{2-p} m' r^{2-|\xi|} (\xi d - 2)^{\xi d-3} \exp[(\xi d - 1) q(x)], \quad x \in \Omega, |\xi| \geq 2, \quad (4.25)$$

and

$$|D^\xi u^{p+1}| \leq 2^{2-p} m', \quad |\xi| \leq 1. \quad (4.26)$$

If  $|\xi + \alpha| \geq 2$  and  $|\alpha| \leq 1$ , then (4.25) gives us

$$\begin{aligned} |D^\xi(D^\alpha u^{p+1})| &\leq 2^{2-p} m' r^{2-|\alpha|} r^{-|\xi|} (\xi d + \alpha d - 2)^{\xi d + \alpha d - 3} \exp((\xi d + \alpha d - 1) q(x)) \\ &\leq 2^{2-p} m' r^{-|\xi|} \xi d^{\xi d-1} \exp[(\xi d + 1) q(x)], \end{aligned}$$

since  $r \leq 1$ ,  $\alpha d \leq 2$ . For  $|\xi + \alpha| \leq 1$  we use (4.26) and get

$$|D^\xi(D^\alpha u^p)| \leq 2^{2-p} m' r^{-|\xi|} \xi d^{\xi d-1} \exp[(\xi d + 1) q(x)].$$

Then we use Lemma 4.1 and get

$$|D^\xi(a_\alpha D^\alpha u^{p+1})| \leq 2^{2-p} m_\alpha m' c r^{-|\xi|} \xi d^{\xi d-1} \exp[(1 + \xi d) q(x)].$$

This and (4.21) tell us that

$$|D^\xi f^{p+1}| \leq \left(c \sum m_\alpha\right) 2^{2-p} m' r^{-|\xi|} \xi d^{\xi d-1} \times \exp[(1 + \xi d) q(x)].$$

Then a look at (4.11) completes the proof of (4.24). So (4.25) and (4.26) are true for all  $p$  too. That implies (4.22) and completes the proof of Theorem 1.5.

## 5. ESTIMATES OF DERIVATIVES

As we mentioned in the Introduction this section has been rewritten. We also take this opportunity to trace the ideas lying behind Lemma 5.1 below and our use of it. The first example of a problem leading to nonanalytical estimates

of Gevrey type seems to be a counterexample by Kovalevskij [E8] showing that the Cauchy problem for the heat equation  $\partial u / \partial t - \partial^2 u / \partial x^2 = 0$  is not always solvable in the class of analytic solutions when the initial datum is given at time  $t = 0$ . Le Roux [E12] and Holmgren [E6] showed that with data in Gevrey classes 2 on  $x = 0$  there exists a solution analytic in the  $x$ -variable. Then Gevrey in [E5] introduced the classes nowadays called Gevrey classes. He also solved the Cauchy problem with data on  $x = 0$  for the heat equation with added "lower" order terms with coefficients in proper Gevrey classes, in which he uses his version of Lemma 5.1. So we may say that he is the one behind it. But he solves the Cauchy problem using an explicit form of the solution of the inhomogeneous heat equation with zero initial data. The first one to use the corresponding idea on successive approximations was Pucci [14] when he solved a general linear Cauchy problem for equations with coefficients in proper Gevrey classes. More transparent versions have been used by Lax [E11] and Friedman [E4] in the proof of different versions of the Cauchy-Kovalevskij theorem. They show that formal power series solutions in the time variable are convergent using estimates of the same type as that in Lemma 5.1 (see also Persson [E13, 10, 11] and Shinbrot and Welland [E16]). Now we give the "canonical" proof of the lemma.

**LEMMA 5.1.** *Let  $d = (d_1, \dots, d_n) \in \mathbb{R}^n$ ,  $d_j \geq 1$ ,  $1 \leq j \leq n$ . Let  $\nu$  and  $\xi$  be multi-indices with nonnegative components. Then there exists a constant  $c$  independent of  $n$ ,  $d$ , and  $\xi$  such that*

$$\sum_{\nu \leq \xi} \binom{\xi}{\nu} (\xi - \nu)^{(\xi - \nu)d - 1} \nu d^{\nu d - 1} \xi d^{1 - \xi d} \leq c. \quad (5.1)$$

Here we have used the natural extension to  $\mathbb{R}^n$  of the notation for  $\mathbb{R}^2$  in the Introduction. We have let

$$\binom{\xi}{\nu} = \prod_{j=1}^n \binom{\xi_j}{\nu_j} \quad \text{and} \quad \nu \leq \xi \Leftrightarrow \nu_j \leq \xi_j, \quad 1 \leq j \leq n.$$

*Remark.* The proof will show that  $c = 4$  is the best constant in (5.1).

*Proof.* Let  $x, y \in \mathbb{R}^n$ . The "binomial" formula gives

$$(x + y)^\xi = \sum_{\nu \leq \xi} \binom{\xi}{\nu} x^\nu y^{\xi - \nu}.$$

Let  $x_1 = \dots = x_n = t$  and let  $y_1 = \dots = y_n = 1$ . We get

$$\sum_{j=0}^{|\xi|} \binom{|\xi|}{j} t^j = (1 + t)^{|\xi|} = \sum_{\nu \leq \xi} \binom{\xi}{\nu} t^{|\nu|}.$$



This implies that

$$\binom{|\xi|}{j} = \sum_{\substack{|\nu|=j \\ \nu \leq \xi}} \binom{\xi}{\nu}. \quad (5.2)$$

We also notice that for  $\nu \leq \xi$ ,  $0 \neq \nu \neq \xi$ ,

$$(\nu d / \xi d)^{\nu d - 1} \leq (|\nu| / |\xi|)^{|\nu| - 1}, \quad (5.3)$$

since  $d_j \geq 1$ ,  $1 \leq j \leq n$ . Let  $|\xi| = k$ . Now (5.2) and (5.3) show that the left member of (5.1) is smaller than

$$A = 2 + \sum_{j=1}^{k-1} \binom{k}{j} (k-j)^{k-j-1} j^{j-1} k^{-k+1}.$$

From [E15, formula (20), p. 20] we get the following identity letting  $x = y = 1$ ,

$$\sum_{j=0}^{k-2} \binom{k-2}{j} (1+j)^{j-1} (k-j-1)^{k-j-2} = 2k^{k-3}.$$

This is equivalent to

$$\sum_{j=1}^{k-1} \binom{k-2}{j-1} j^{j-2} (k-j)^{k-j-2} = 2k^{k-3},$$

or

$$(k(k-1))^{-1} \sum_{j=1}^{k-1} \binom{k}{j} j^{j-1} (k-j)^{k-j-1} = 2k^{k-3}.$$

In other words

$$\sum_{j=1}^{k-1} \binom{k}{j} j^{j-1} (k-j)^{k-j-1} k^{-k+1} = 2 - 2/k,$$

for all  $k \geq 2$ . The lemma is proved.

*Proof of Lemma 4.1.* Let  $f$  and  $g$  satisfy (4.3) and (4.4), respectively. We use Leibniz's formula on  $D^\xi(fg)$ , then the estimates (4.3) and (4.4), and at last Lemma 5 gives

$$\begin{aligned} |D^\xi(fg)| &\leq \sum_{\nu \leq \xi} \binom{\xi}{\nu} |D^{\xi-\nu}g| |D^\nu f| \\ &\leq \sum_{\nu \leq \xi} \binom{\xi}{\nu} m r^{-|\xi-\nu|} (\xi-\nu)^{(\xi-\nu)d-1} \nu^{\nu d-1} m' r^{-|\nu|} (\nu d)^{\nu d-1} \\ &\quad \times \exp[(1+\nu d)q(x)] \leq m m' c r^{-|\xi|} \xi d^{\xi d-1} \exp[(1+\xi d)q(x)]. \end{aligned}$$

We have also noticed that  $e^j$  is increasing. The lemma is proved.

The special case with  $q = 0$  shows us that the Gevrey classes with  $d \geq (1, \dots, 1)$  are closed under multiplication. An easy argument shows that they are also closed under differentiation. Here we must adjust the  $r$  of the estimate, not only choose a new constant  $m$  in our estimate. We have used these two facts when we derived (4.23).

## 6. THE HARD PART

We now prove Lemma 4.3. The first statement of the lemma is easy to verify so we concentrate upon the second one. In this section we write  $\xi = (i, j)$ , instead of  $\xi = (\xi_1, \xi_2)$ . We also define

$$A(k) = k^{k-1}, \quad k \geq 1, A(k) = 1, 0 \leq k \leq 1, \quad (6.1)$$

and

$$E(k, t) = \exp[(1 + k) e^2 |t|/r], \quad k \geq 0. \quad (6.2)$$

We let  $d$  denote a number here. More specifically we let  $d = d_2 \geq 1$ . The number  $k$  in (6.1) and (6.2) is of the form  $k = jd$ , with  $j \geq 0$  an integer. In these numbers  $k$  both  $A(k)$  and  $E(k, t)$  are nondecreasing. In this notation Lemma 5.1 takes the form

$$\sum_{j=0}^k \binom{k}{j} A((k-j)d) A(jd) \leq cA(kd), \quad \text{all } k. \quad (6.3)$$

We also notice that

$$A(k) \leq A(k-p) k^p e^p, \quad p \leq k, \quad (6.4)$$

and if  $e^2 |t|/r \leq 1$ , then

$$E(k, t) \leq E(k-p, t) e^p. \quad (6.5)$$

We leave the proofs to the reader.

*Proof of Lemma 4.3.* The first step in the proof is to perform all differentiations in the expression of the left member of (4.9) and (4.10). To facilitate the bookkeeping of the terms that arise we write the resulting expression in the following form

$$D_1^i D_2^j \int_0^{x_2} \int_0^t g(x_1 - \varphi(x_2) + \varphi(t), s) ds dt = B_{i,j}^1 + B_{i,j}^2 + B_{i,j}^3, \quad (6.6)$$

where  $B_{i,j}^1$  denotes the sum of all terms that contain a double integral,  $B_{i,j}^2$  the sum of those containing a single integral, and  $B_{i,j}^3$  the sum of those without integral signs. We define

$$Q_{j,j} = b^j, \quad j \geq 0, \quad (6.7)$$

$$Q_{j,0} = 0, \quad j > 0, \quad (6.8)$$

and

$$Q_{j+1,k} = bQ_{j,k-1} + D_2 Q_{j,k}, \quad 0 < k \leq j. \quad (6.9)$$

We remember that  $\varphi' = b$ . A straightforward calculation shows that

$$B_{i,j}^1 = \sum_{k=0}^j Q_{j,k} \int_0^{x_2} \int_0^t (D_1^{i+k} g)(x_1 - \varphi(x_2) + \varphi(t), s) ds dt, \quad (6.10)$$

$$B_{i,j}^2 = \sum_{l=0}^{j-1} \sum_{k=0}^l \left( \int_0^{x_2} (D_1^{i+k} g)(x_1, s) ds \right) D_2^{j-1-l} Q_{l,k}, \quad (6.11)$$

and

$$B_{i,j}^3 = \sum_{p=1}^{j-1} \sum_{l=0}^{p-1} \sum_{k=0}^l D_2^{j-1-p} [(D_1^{i+k} g)(x_1, x_2) D_2^{p-1-l} Q_{l,k}]. \quad (6.12)$$

*Remark.* If  $j \geq 1$  the sum in (6.10) actually starts at  $k = 1$  since  $Q_{j,0} = 0$  then. Similarly if  $j \geq 2$  the sums in (6.11) start at  $k = 1, l = 1$  since  $D_2^{j-1-l} Q_{l,k} = 0$  if  $(l, k) = (0, 0)$  or  $(l, k) = (1, 0)$ . For  $j = 1$  there is just one term in (6.11) and none for  $j = 0$ . The expression of  $B_{i,j}^3$  in (6.12) is empty if  $j \leq 1$ . For  $j \geq 2$  it contains just one term with  $l = k = 0$ , namely,  $D_1^i D_2^{j-2} g$ . If that term is taken out the rest of the sum can be taken from  $k = 1$  and  $l = 1$ , too, and the first summation starts at  $p = 2$ .

We now assert that

$$|D_2^s Q_{l,k}| \leq m(mc)^{k-1} \binom{l-1}{k-1} R^{-(s+l-k)} A((s+l-k)d),$$

$$0 < k \leq l, \quad s \geq 0, \quad |x_2| < \rho, \quad (6.13)$$

where  $c$  is defined in Lemma 4.1, and  $\rho, m$ , and  $R$  are defined in Lemma 4.3.

If  $k = l$  then we conclude from (6.7), (4.8), and Corollary 4.2 that (6.13) is true for  $s \geq 0$ . Notice that  $d = d_2$  here. Finally assume that (6.13) is true for a certain  $l, 0 < k \leq l$  and all  $s$ . This is certainly the case for  $l = 1$ . Now take  $0 < k \leq l$ . We use (6.9), Leibniz' formula, (6.13), and (4.8). For  $k > 1$  we get

$$\begin{aligned} & |D_2^s Q_{l+1,k}| \\ & \leq |D_2^s (bQ_{l,k-1})| + |D_2^{s+1} Q_{l,k}| \\ & \leq \sum_{t=0}^s \binom{s}{t} |D_2^{s-t} b| |D_2^t Q_{l,k-1}| + |D_2^{s+1} Q_{l,k}| \\ & \leq \sum_{t=0}^s \binom{s}{t} m^2 (mc)^{k-2} R^{-(s+l-k+1)} \binom{l-1}{k-2} A((s-t)d) A((t+l-k+1)d) \\ & \quad + m(mc)^{k-1} \binom{l-1}{k-1} R^{-(s+1+l-k)} A((s+1+l-k)d). \end{aligned}$$

Now we notice that

$$A((t + l - k + 1)d)(A((s + l - k + 1)d))^{-1} \leq A(td)(A(sd))^{-1}.$$

We use this, (6.3), and  $\binom{l-1}{k-2} + \binom{l-1}{k-1} = \binom{l}{k-1}$ . We get

$$|D_2^s Q_{l+1,k}| \leq m(mc)^{k-1} R^{-(s+l+1-k)} \binom{l}{k-1} A((s + l - k + 1)d).$$

By that we have proved that (6.13) is true when  $l$  is replaced by  $l + 1$  and if  $1 < k \leq l$ . We notice that  $D_2^s Q_{l+1,1} = D_2^{s+1} Q_{l,1}$ . So (6.13) is also true when  $k = 1$  and  $l$  is replaced by  $l + 1$ . The case  $Q_{l+1,l+1}$  is treated earlier. So (6.13) is always true.

We can now deduce estimates for the terms  $B_{i,j}^k$ ,  $k = 1, 2, 3$ , using (6.13) and the assumption (4.4) which in our present notation reads

$$|D_1^i D_2^j g| \leq m' r^{-(i+j)} A((i+j)d) E((i+j)d, x_2), \quad x \in \Omega. \quad (6.14)$$

However, it is a rather complicated task to reduce the resulting estimates into a manageable form so we have to present the computations in detail. We look at  $B_{i,j}^1$  first, (6.10). Let  $j \geq 1$ .

$$\begin{aligned} |B_{i,j}^1| &\leq \sum_{k=1}^j m' r^{-i-k} A((i+k)d) m(mc)^{k-1} \binom{j-1}{k-1} \\ &\quad \times R^{-(j-k)} A((j-k)d) \left| \int_0^{\omega_2} \int_0^t E((i+j)d, s) ds dt \right|. \end{aligned}$$

We have used  $E((i+k)d, s) \leq E((i+j)d, s)$  here. From (4.13) we get  $mc < \frac{1}{2}$ ,  $r < R$ . We notice that (6.3) gives

$$\begin{aligned} &\sum_{k=1}^j \binom{j-1}{k-1} A((i+k)d) A((j-k)d) \\ &\leq \sum_{k=1}^j \binom{i+j}{i+k} A((i+k)d) A((j-k)d) \leq c A((i+j)d). \end{aligned}$$

This together with integration of the double integral gives us

$$|B_{i,j}^1| \leq m' r^{-i-j} A((i+j)d) (1 + (i+j)d)^{-2} r^2 e^{-3} \times E((i+j)d, x_2). \quad (6.15)$$

It is clear that (6.15) is also true for  $j = 0$ . Now we use (6.4) and (6.5) with  $p = 2$  for the case  $i + j \geq 2$ . We get

$$|B_{i,j}^1| \leq m r^{-i-j+2} A((i+j)d - 2) E((i+j)d - 2, x_2), \quad i + j \geq 2. \quad (6.16)$$

When  $i + j \leq 1$  a short computation gives

$$|B_{i,j}^1| \leq m', \quad i + j \leq 1, \quad (6.17)$$

since  $d \leq 2$ .

The estimate of  $B_{i,j}^2$  is obtained in much the same way. We estimate the right member of (6.11) using (6.13) and (6.14). Then we replace  $E((i + k)d, s)$  by  $E((i + j - 1)d, s)$  before we integrate. For  $j \geq 2$  we get

$$\begin{aligned} |B_{i,j}^2| &\leq \sum_{l=1}^{j-1} \sum_{k=1}^l m' m [1 + (i + j - 1)d]^{-1} r e^{-2} E((i + j - 1)d, x_2) \\ &\quad \times r^{-(i+k)} (mc)^{k-1} \binom{l-1}{k-1} R^{-(j-1-k)} A((i + k)d) A((j - k - 1)d). \end{aligned}$$

We interchange the summations. We use  $mc < \frac{1}{2}$ ,  $r < R$  and get

$$\begin{aligned} |B_{i,j}^2| &\leq mm' r^{-(i+j)+2} E((i + j - 1)d, x_2) e^{-2} [1 + (i + j - 1)d]^{-1} \\ &\quad \times \sum_{k=1}^{j-1} A((i + k)d) A((j - k - 1)d) \sum_{l=k}^{j-1} \binom{l-1}{k-1}. \end{aligned}$$

We notice that

$$\sum_{l=k}^{j-1} \binom{l-1}{k-1} = \binom{j-1}{k}. \quad (6.18)$$

We then argue as in the last step in the estimation of  $B_{i,j}^1$ . We get

$$|B_{i,j}^2| \leq m' r^{-(i+j)+2} E((i + j - 1)d, x_2) e^{-2} (1 + (i + j - 1)d)^{-1} \times A((i + j - 1)d)$$

This estimate also holds for  $j = 1$ . For  $i + j \geq 2$  we use (6.4) and (6.5) with  $p = 1$ . We notice that  $d \geq 1$ . We get

$$|B_{i,j}^2| \leq m' r^{-(i+j)+2} E((i + j)d - 2, x_2) A((i + j)d - 2). \quad (6.19)$$

For  $i + j \leq 1$  we only have

$$|B_{0,1}^2| \leq m'. \quad (6.20)$$

The estimation of  $B_{i,j}^3$  is more complicated. In our estimate we use that  $4r \leq R$ , not just that  $r \leq R$ . As mentioned before,  $B_{i,j}^3 = 0$  for  $j < 2$  and  $B_{i,2}^3 = D_1^i g$  which we can estimate by (6.14). We also recall that  $D_1^i D_2^{j-2} g$  is the only non-vanishing term in (6.12) with  $l = 0$  or  $k = 0$ . We use this and Leibniz' formula to rewrite (6.12) as

$$\begin{aligned} B_{i,j}^3 - D_1^i D_2^{j-2} g &= \sum_{p=2}^{j-1} \sum_{l=1}^{p-1} \sum_{k=1}^l \sum_{q=0}^{j-1-p} \binom{j-1-p}{q} (D_1^{i+k} D_2^q g) \\ &\quad \times D_2^{j-2-l-q} Q_{l,k}, \quad j \geq 3. \end{aligned}$$

We use (6.13) and (6.14). Since  $q + k \leq j - 2$  we may replace  $E((i + k + q)d, x_2)$  by  $E((i + j - 2)d, x_2)$  in the estimate of each term. For  $j \geq 3$  we get

$$\begin{aligned} & |B_{i,j}^3 - D_1^i D_2^{j-2} g| \\ & \leq mm' r^{-i-j+2} E((i + j - 2)d, x_2) \\ & \quad \times \sum_{p=2}^{j-1} \sum_{l=1}^{p-1} \sum_{k=1}^l \sum_{q=0}^{j-1-p} \binom{j-1-p}{q} \binom{l-1}{k-1} (mc)^{k-1} (r/R)^{j-2-k-q} \\ & \quad \times A((i + k + q)d) A((j - 2 - k - q)d). \end{aligned}$$

Just as in the derivation of (6.19), we interchange the  $l$  and  $k$  summations, and get

$$\begin{aligned} & |B_{i,j}^3 - D_1^i D_2^{j-2} g| \\ & \leq mm' r^{-(i+j-2)} E((i + j - 2)d, x_2) \\ & \quad \times \sum_{p=2}^{j-1} \sum_{k=1}^{p-1} \sum_{q=0}^{j-1-p} \binom{j-1-p}{q} \binom{p-1}{k} (mc)^{k-1} (r/R)^{j-2-k-q} \\ & \quad \times A((i + k + q)d) A((j - 2 - k - q)d) \end{aligned} \quad (6.21)$$

We shall need the following lemma.

**LEMMA 6.1.** *Let  $j \geq 3$ ,  $p, q, i \geq 0$ , be integers such that  $0 \leq q \leq j - 1 - p$ ,  $2 \leq p \leq j - 1$ ,  $1 \leq k \leq p - 1$ . Let  $d \geq 1$ . Then we have*

$$\begin{aligned} & A((i + k + q)d) A((j - 2 - k - q)d) (A((i + j - 2)d))^{-1} \\ & \leq A(q) A(j - p - 1 - q) (A(j - p - 1))^{-1}. \end{aligned}$$

*Proof.* First we notice that

$$\begin{aligned} & A((i + k + q)d) A((j - 2 - k - q)d) (A((i + j - 2)d))^{-1} \\ & \leq A(k + q) A(j - 2 - k - q) (A(j - 2))^{-1}. \end{aligned}$$

Then we notice that

$$k \rightarrow A(k + q) A(j - 2 - q - k)$$

is a convex function. Thus we have

$$\begin{aligned} & A(k + q) A(j - 2 - k - q) \\ & \leq \max(A(q) A(j - 2 - q), A(q + p - 1) A(j - 1 - p - q)). \end{aligned}$$

We also notice that

$$A(q) A(j - 2 - q) (A(j - 2))^{-1} \leq A(q) A(j - p - 1 - q) (A(j - p - 1))^{-1},$$

and

$$\begin{aligned} & A(q+p-1) A(j-1-p-q)(A(j-2))^{-1} \\ & \leq A(q) A(j-p-1-q)(A(j-p-1))^{-1}. \end{aligned}$$

The lemma is proved.

Lemma 6.1 applied to (6.21) gives us

$$\begin{aligned} & |B_{i,j}^3 - D_1^i D_2^{j-2} g| \\ & \leq m' r^{-i-j+2} E((i+j-2)d, x_2) \\ & \quad \times A((i+j-2)d) \sum_{p=2}^{j-1} \sum_{q=0}^{j-1-p} \binom{j-1-p}{q} (r/R)^{j-2-q-p-1} \\ & \quad \times A(q) A(j-1-p-q)(A(j-1-p))^{-1} c^{-1} \\ & \quad \times \sum_{k=1}^{p-1} \binom{p-1}{k} (mc)^k (r/R)^{p-1-k}. \end{aligned} \quad (6.22)$$

We notice that  $j-2-q-p-1 \geq 0$ ,  $r/R \leq 1$ . We also notice that

$$\sum_{k=1}^{p-1} \binom{p-1}{k} (mc)^k (r/R)^{p-1-k} = (mc + r/R)^{p-1} - (r/R)^{p-1},$$

and that because of (4.13)

$$\sum_{p=2}^{j-1} (mc + r/R)^{p-1} \leq (mc + r/R)(1 - mc - r/R)^{-1} \leq 1.$$

These facts together with  $d \geq 1$ , (6.22), and (6.3) show that for  $j \geq 3$

$$|B_{i,j}^3 - D_1^i D_2^{j-2} g| \leq m' r^{(-i-j-2)} A((i+j)d-2) E((i+j)d-2, x_2).$$

This and (6.14) show that

$$|B_{i,j}^3| \leq 2m' r^{-i-j-2} A((i+j)d-2) E((i+j)d-2, x_2), \quad i \geq 0, \quad j \geq 2. \quad (6.23)$$

Now (6.16), (6.19), and (6.23) give (4.9). Finally, (6.17) and (6.20) give (4.10). Lemma 4.3 is proved. By this the full proof of Theorem 1.5 is also completed.

## REFERENCES

1. M. S. BAOUENDI AND C. GOULAOUIC, Cauchy problems with characteristic initial hypersurface, *Comm. Pure Appl. Math.* **26** (1973), 455–475.
2. M. D. BRONŠTEIN, On Differential equations having non-analytic solutions, *Uspehi Mat. Nauk* **30** (1975), 227–228, (Russian).

3. L. HÖRMANDER, On the theory of general partial differential operators, *Acta Math.* **94** (1955), 161–248.
4. L. HÖRMANDER, Non-uniqueness for the Cauchy problem, in “Fourier integral operators and partial differential equations” (Coll. Int., Nice, 1974), Lecture Notes in Mathematics, No. 459, Springer, Berlin/Heidelberg/New York, 1975.
6. H. KOMATSU, Irregularity of characteristic elements and construction of null-solutions, *J. Fac. Sci. Univ. Tokyo Sect. IA* **23** (1976), 297–342.
7. J. PERSSON, Exponential majorization and global Goursat problems, *Math. Ann.* **178** (1978), 271–276.
8. J. PERSSON, On the supports of solutions of linear partial differential equations with analytic coefficients, *Ann. Mat. Pura Appl. (4)* **91** (1972), 79–86.
9. J. PERSSON, On uniqueness cones, velocity cones and  $P$ -convexity, *Ann. Mat. Pura Appl. (4)* **96** (1973), 69–87.
10. J. PERSSON, Non-uniqueness in the characteristic Cauchy problem when the coefficients are analytic, *Mathematicheskoe* **27** (1972), 145–152.
11. J. PERSSON, Semi-global null solutions and  $P$ -convexity, *Boll. Un. Mat. Ital. (4)* **8** (1973), 20–28.
12. J. PERSSON, Non-uniqueness of Täcklind type in the characteristic Cauchy problem, *Matematicheskoe* **29** (1974), 162–175.
13. J. PERSSON, Correction of “Semi-global null solutions and  $P$ -convexity”, *Boll. Un. Mat. Ital. (4)* **11** (1975), 518–524.
14. C. PUCCI, Nuove ricerche sul problema di Cauchy, *Mem. Accad. Sci. Torino, Cl. Sci. Fis. Mat. Natur.* **1** (1954), 45–67.
15. L. SCHWARTZ, “Théorie des distributions,” Tome I. Hermann, Paris, 1950.
16. S. TÄCKLIND, Sur les classes quasianalytiques des solutions des équations aux dérivées partielles du type parabolique, *Nova Acta Soc. Sci. Upsalensis (4)* **10** (1936), 1–57.
17. F. TREVES, “Linear Partial Differential Equations with Constant Coefficients,” Gordon & Breach, New York/London/Paris, 1966.
18. F. TREVES, Discrete phenomena in uniqueness in the Cauchy problem, *Proc. Amer. Math. Soc.* **46** (1974), 229–233.
19. A. TYHONOV, Théorèmes d’unicité pour l’équation de la chaleur, *Mat. Sb.* **42** (1935), 199–216.

## EXTRA REFERENCES

- E1. R. BADER ET P. GERMAIN, Solutions élémentaires de certaines équations aux dérivées partielles du type mixte, *Bull. Soc. Math. France* **81** (1953), 145–174.
- E2. I. S. BEREZIN, On Cauchy’s problem for linear equations of the second order with initial conditions on a parabolic line, *Mat. Sb.* **24** (1949), 301–320.
- E3. R. CONTI, Sul problema di Cauchy per l’equazione  $y^{2\alpha}k^2(x, y) z_{xx} - z_{yy} = f(x, y, z, z_x, z_y)$ , con i dati sulla linea parabolica, *Ann. Mat. Pura Appl.* **31** (1950), 303–326.
- E4. A. FRIEDMAN, A new proof and generalizations of the Cauchy-Kowalewski theorem, *Trans. Amer. Math. Soc.* **98** (1961), 1–20.
- E5. M. GEVREY, Sur la nature analytique des solutions des équations aux dérivées partielles, *Ann. École Norm. Sup.* **35** (1918), 127–190.
- E6. E. HOLMGREN, Sur l’équation de la propagation de la chaleur, *Ark. Mat. Fys. Astr.* **4** (1908), 14, 18.
- E7. V. JA. IVRIĬ, Correctness of the Cauchy problem in Gevrey classes for non-strictly hyperbolic operators, *Mat. Sb.* **96** (138), (1975), 390–413; English transl., *Math. USSR-Sb* **25** (1975), 365–387.



- E8. S. KOWALEWSKI, Zur Theorie der partiellen Differentialgleichungen, *J. Reine Angew. Math.* **80** (1875), 1–32.
- E9. E. KULIKOV, Cauchy's problem for hyperbolic equations of degenerate type on the initial line, *Differencial'nye Uravenija* **11** (1975), 512–522; English transl., *Differential Equations* **11** (1975), 385–393.
- E10. N. A. LAR'KIN, Cauchy's problem for a second order quasi-linear hyperbolic equation with initial data on a line of parabolicity, *Differencial'nye Uravenija* **8** (1972), 76–84; English transl., *Differential Equations* **8** (1972), 57–63.
- E11. P. LAX, Nonlinear hyperbolic equations, *Comm. Pure Appl. Math.* **4** (1953), 231–258.
- E12. J. LE ROUX, Sur les intégrales analytiques de l'équation  $\partial^2 u / \partial y^2 = \partial u / \partial x$ , *Bull. Sci. Math.* **19** (1895), 127–129.
- E13. J. PERSSON, New proofs and generalization of two theorems by Lednev for Goursat problems, *Math. Ann.* **178** (1968), 184–208.
- E14. M. H. PROTTER, The Cauchy problem for a hyperbolic second order equation with data on the parabolic line, *Canad. J. Math.* **6** (1954), 542–553.
- E15. J. RIORDAN, "Combinatorial Identities," Wiley, New York/London/Sidney, 1968.
- E16. M. SHINBROT AND R. R. WELLAND, The Cauchy–Kowalewskaya theorem, *J. Math. Anal. Appl.* **55** (1976), 757–772.